

The correlation energy as an explicit functional of the one-particle density matrix from a determinantal reference state

James P. Finley*

*Department of Physical Sciences, Eastern New Mexico University,
Station #33, Portales, NM 88130 and*

*Department of Applied Chemistry, Graduate School of Engineering,
The University of Tokyo, 7-3-1 Hongo,
Bunkyo-ku, Tokyo, 113-8656 Japan*

(Dated: February 2, 2008)

Abstract

Using an approach based on many body perturbation theory, the correlation energy $\mathcal{E}_{\text{co}}$ is expressed as an explicit functional of ρ_1 , v , and v_s , where ρ_1 is the one-particle density matrix from the noninteracting, or reference, determinantal-state; v is the external potential from the interacting, or target, state; v_s is the (kernel of the) external potential from the noninteracting determinantal-state. In other words we have $\mathcal{E}_{\text{co}}[\rho_1, v, v_s]$. Another possibility is the following explicit functional: $\mathcal{E}_{\text{co}}[\rho_1, v_{\text{co}}, v_s]$, where v_{co} is the (kernel of the) correlation potential from the noninteracting Hamiltonian. The proposed method can, in principle, be used to compute $\mathcal{E}_{\text{co}}$ in a very accurate and efficient manner, since, like the Kohn–Sham approach, there are no virtual orbitals to consider. However, in contrast to the Kohn–Sham approach, $\mathcal{E}_{\text{co}}$ is a known, explicit functional that can be approximated in a systematic manner. For simplicity, we only consider noninteracting closed-shell states and target states that are nondegenerate, singlet ground-states; so, in that case, ρ_1 denotes the spin-less one-particle density matrix from the determinantal reference state.

I. INTRODUCTION

The Kohn-Sham version of density functional theory has been very successful in the description of electronic structure for quantum chemistry and condensed matter physics.^{1,2,3,4,5,6,7} Unlike pure density functional approaches,^{1,2,8} the Kohn-Sham method uses a set of occupied orbitals from a noninteracting state, where this determinantal state shares its electron density ρ with the target, or interacting, state $|\Psi\rangle$. In the Kohn-Sham approach, the kinetic energy (through the) first order is not an explicit functional of ρ , but, instead, this functional depends on the one-particle density matrix ρ_1 from the noninteracting, determinantal state. The Kohn-Sham approach requires as input the exchange-correlation functional E_{xc}^{KS} —or equivalently the exchange E_x and correlation-energy E_{co}^{KS} functionals—where E_{xc}^{KS} is required to be an explicit functional of ρ . Unfortunately, E_{xc}^{KS} is an unknown functional, and there is no systematic method to improve approximations.

The optimized potential method^{9,10,11,12,13,14,15} is a density functional approach that can convert a nonlocal operator into a local one, where the exchange-correlation functionals can depend on both the occupied and virtual orbitals. Unfortunately, this method lacks the efficiency of other Kohn-Sham approaches. Furthermore, the optimized potential method introduces functional that—in contrast to many wave function methods—are not invariant to a unitary transformation of either the occupied or virtual orbitals; the local potentials also depend on the orbital energies.

High levels of approximations—beyond the Kohn-Sham approaches—can be obtained by wave function methods,^{16,17,18,19} including the coupled cluster method, many-body perturbation theory, and configuration interaction. Often, however, these methods are much less efficient than the Kohn Sham approaches, where, typically, wave function methods consider a large number of 2-electron molecular integrals, depending on both the occupied and virtual orbitals, and these integrals must be computed and utilized in calculations involving large atomic-orbital basis sets.

In order to improve the efficiency of the wave function methods, especially in regards to their scaling with molecular size, perturbative methods have been developed based on localized molecular orbitals.^{20,21,22,23,24,25,26} An alternative approach uses a Laplace transform to remove the energy denominators in perturbation theory, yielding approaches involving correlation energy expressions that depend explicitly on the atomic-orbital basis set.^{27,28}

This method has also been used with the coupled cluster theory.²⁹

Our research interest is in the development of methods that bridge the gap between wave function approaches and Kohn–Sham density functional theory. However, these two different approaches already share some common features. For example, as mentioned above, Kohn–Sham introduces a determinantal wave function and treats the first-order kinetic energy in the same manner as in wave function methods, i.e., as a functional of the one-body density matrix from a determinant. Furthermore, hybrid density functionals,^{30,31,32,33} including B3LYP,^{30,34} introduce a component of the exact exchange—a functional of ρ_1 —even though these approaches violate the Hohenberg–Kohn theorem⁸ by using the nonlocal exchange operator.

A simple generalization of the Kohn–Sham functionals involves using the exact exchange energy $\mathcal{E}_x$ (with its corresponding non-local operator) and, in addition, a correlation-energy functional that also depends on ρ_1 . Such an approach could probably be derived within a variant of the Kohn–Sham scheme. However, it is also reasonable to base such a formalism on wave function methods, since, for example, the correlation-energy, say $\mathcal{E}_{\text{co}}$, from wave function methods is an implicit functional of ρ_1 , and does not implicitly depend on the orbitals, even though there is an explicit dependence. (This is easily proven by noting that we have $\mathcal{E}_{\text{co}} = \mathcal{E} - E_1[\rho_1]$, where $\mathcal{E}$ is the exact electron energy, and E_1 is the first order energy that is determined by ρ_1 .) Furthermore, such a formalism can be based on any reasonable orbitals: Hartree–Fock, Brueckner, and natural, but there is probably no advantage to using Kohn–Sham orbitals. In addition, however, when considering a wave function approach, $\mathcal{E}_{\text{co}}$ also has a natural dependence on the external potential v from the interacting or target state $|\Psi\rangle$. (This is easily proven from coupled cluster theory by noting the Fock operator $\hat{F}$ appearance in the coupled equations and noting that $\hat{F}$ depends on v .) Therefore, it is reasonable to have $\mathcal{E}_{\text{co}}$ depending on both on v and ρ_1 . Since v is a known function defining the molecular structure, the inclusion of the v should not produce any difficulties.

Below we introduce an approach based on many body perturbation theory,^{16,17,18,35} where the correlation energy $\mathcal{E}_{\text{co}}$ is expressed as an explicit functional of ρ_1 , v , and v_s , where ρ_1 is the one-particle density matrix from the noninteracting, or reference, state; v is the external potential from the interacting state $|\Psi\rangle$; v_s is the (kernel of the) external potential from the noninteracting state. In other words we have $\mathcal{E}_{\text{co}}[\rho_1, v, v_s]$. Here, the v_s explicit dependence appears because the individual diagrams from the perturbative expansion depend on

the zeroth-order Hamiltonian, or, equivalently, on v_s , since—together with the number of electrons— v_s defines the zeroth-order Hamiltonian. However, unless the perturbation expansion is truncated at some order, it is easily proven that $\mathcal{E}_{\text{co}}$ does not implicitly depend on v_s , since an exact $\mathcal{E}_{\text{co}}$ does not depend of the zeroth-order Hamiltonian. (Infinite order summations, e.g., the coupled cluster method, also have $\mathcal{E}_{\text{co}}$ not depending on v_s .)

This method presented below can, in principle, be used to compute $\mathcal{E}_{\text{co}}$ in an accurate and efficient manner, since, like the Kohn–Sham approach, there are no virtual orbitals to consider. However, in contrast to the Kohn–Sham approach, $\mathcal{E}_{\text{co}}$ is a known, explicit functional that can be approximated in a systematic manner. Furthermore, in contrast to the optimized potential method,^{9,10,11,12,13,14,15} the functionals are invariant to a unitary transformation of occupied orbitals. The proposed method, like the efficient wave function methods mentioned above,^{27,28,29} removes the energy denominators by using Laplace transforms, as well as other techniques, to obtain the correlation energy as an explicit functional of ρ_1 , v , and v_s . For simplicity, we only consider noninteracting closed-shell states and target states that are nondegenerate, singlet ground-states; so, in that case, ρ_1 denotes the spin-less one-particle density matrix from the determinantal reference state.

Let us also mention that we have recently derived a generalization of the Kohn–Sham approach in which the correlation energy $\mathcal{E}_{\text{co}}$ is assumed to be an explicit functional of v and ρ_1 .^{36,37,38,39} In a similar Brueckner-orbital method, called reference-state one-particle density matrix theory,^{40,41,42} $\mathcal{E}_{\text{co}}$ is also assumed to be an explicit functional of v and ρ_1 , where for closed-shell systems, the local density approximation (LDA) and the LYP, BLYP, and B3LYP functionals was shown to be compatible with the method.⁴² In both of these two approach, errors from Coulomb self-interactions do not occur, nor the need to introduce functionals defined by a constraint search. By utilizing the method presented here, explicit forms of these two correlation-energy functionals can be obtained.⁴³

II. THE NONINTERACTING STATE

A determinantal state is a wave function that is an eigenfunction of a noninteracting Hamiltonian, say $\hat{H}_{Nv_s}$, where $\hat{H}_{Nv_s}$ has no electron-electron interaction terms. These non-interacting Hamiltonian $\hat{H}_{Nv_s}$ are completely specified by the number of electrons N and

the kernel of an external operator, say v_s . Explicitly we have

$$\hat{H}_{v_s} = \sum_{i=1}^N \hat{\mathcal{F}}_{v_s}(\mathbf{r}_i), \quad (1)$$

where

$$\hat{\mathcal{F}}_{v_s} = -\frac{1}{2}\nabla^2 + \hat{v}_s, \quad (2)$$

and the nonlocal operator $\hat{v}_s$ —which we require to be spin-free—is defined by its kernel v_s , i.e., we have

$$\hat{v}_s\chi(\mathbf{r}_1) = \int d\mathbf{r}_2 v_s(\mathbf{r}_1, \mathbf{r}_2)\chi(\mathbf{r}_2). \quad (3)$$

Using second quantization, our noninteracting Hamiltonian, given by Eq. (1), becomes

$$\hat{H}_{v_s} = \sum_{ij} (i|\hat{\mathcal{F}}_{v_s}|j) \hat{E}_{ij}, \quad (4)$$

where the spin-adapted excitation operator is given by

$$\hat{E}_{ij} = \sum_{\sigma} a_{i\sigma}^{\dagger} a_{j\sigma}, \quad (5)$$

and the one-electrons integrals are spin-free and are written in chemist's notation;¹⁸ these integrals use a spatial orbital set, say $\{\chi\}$; this set has the following form:

$$\psi_{i\sigma}(\mathbf{r}, \omega) = \chi_i(\mathbf{r})\sigma(\omega); \quad \sigma = \alpha, \beta, \quad (6)$$

where the spatial and spin coordinates, $\mathbf{r}$ and ω , can be denoted collectively by $\mathbf{x}$, e.g.,

$$\psi_{i\sigma}(\mathbf{x}) = \psi_{i\sigma}(\mathbf{r}, \omega). \quad (7)$$

It is well known that there is a one-to-one mapping between determinantal states and their one-particle density matrices,^{2,44} say γ . For a closed-shell state we have

$$\gamma(\mathbf{x}_1, \mathbf{x}_2) = \sum_{x \in \{\chi_o\}} \sum_{\sigma} \chi_x(\mathbf{r}_1)\sigma(\omega_1) \chi_x^*(\mathbf{r}_2)\sigma^*(\omega_2) = \sum_{x \in \{\chi_o\}} \chi_x(\mathbf{r}_1)\chi_x^*(\mathbf{r}_2)\delta_{\omega_1\omega_2}, \quad (8)$$

and the spin-less one-particle density-matrix is

$$\rho_1(\mathbf{r}_1, \mathbf{r}_2) = \sum_{\omega} \gamma(\mathbf{r}_1, \omega, \mathbf{r}_2, \omega) = 2 \sum_{x \in \{\chi_o\}} \chi_x(\mathbf{r}_1)\chi_x^*(\mathbf{r}_2), \quad (9)$$

where the sum is over the occupied orbitals from Eq. (6); this set of orbitals is denoted by $\{\chi_o\}$. Comparing Eqs. (8) and (9) yields

$$\gamma(\mathbf{x}_1, \mathbf{x}_2) = \frac{1}{2}\rho_1(\mathbf{r}_1, \mathbf{r}_2)\delta_{\omega_1\omega_2}. \quad (10)$$

Since our closed-shell determinantal states are determined by ρ_1 , henceforth, these kets are denoted by $|\rho_1\rangle$.

Using Eq. (4) and the determinantal state $|\rho_1\rangle$, our eigenvalue problem can be written as

$$\hat{H}_{v_s}|\rho_1\rangle = 2 \left(\sum_{w \in \{\chi_o\}} \epsilon_w \right) |\rho_1\rangle = \mathcal{E}_0 |\rho_1\rangle, \quad (11)$$

where the set of occupied orbitals $\{\chi_o\}$ is defined by the lowest N eigenstates of $\hat{\mathcal{F}}_{v_s}$,

$$\hat{\mathcal{F}}_{v_s} \chi_w = \epsilon_w \chi_w, \quad \chi_w \in \{\chi_o\} \longrightarrow \rho_1, \quad (12)$$

and the right side indicates that these orbitals, from the set $\{\chi_o\}$, determine ρ_1 . The unoccupied, or virtual, orbital set $\{\chi_u\}$ are also obtained from $\hat{\mathcal{F}}_{v_s}$:

$$\hat{\mathcal{F}}_{v_s} \chi_r = \epsilon_r \chi_r, \quad \chi_r \in \{\chi_u\} \longrightarrow \rho_1, \quad (13)$$

where these orbitals also determine ρ_1 , since, for a complete set of orbitals, the unoccupied set determines the occupied set; two sets of orbitals that differ by a unitary transformation are considered equivalent.

While there is no restriction placed upon v_s , normally it contains the external potential v from the interacting state $|\Psi_{Nv}\rangle$, the Coulomb potential $\hat{J}$, and an additional term, the exchange-correlation potential $\hat{v}_{xc}$, that we also permit to depend on the one-particle density matrix:

$$\hat{v}_s = v + \hat{J}_{\rho_s} + \hat{v}_{xc}^{\rho_1} = v + \hat{J}_{\rho_s} + \hat{K}_{\rho_1} + \hat{v}_{co}^{\rho_1}, \quad (14)$$

and the latter expression defines the correlation potential $\hat{v}_{co}^{\rho_1}$, where $\hat{K}$ is the exchange operator, defined below; the Coulomb potential $\hat{J}$, given by

$$\hat{J}_{\rho_s}(\mathbf{r}_1) = \int r_{12}^{-1} \rho_s(\mathbf{r}_2) d\mathbf{r}_2, \quad (15)$$

depends on the electron density ρ_s of the noninteracting state $|\rho_1\rangle$:

$$\rho_s(\mathbf{r}) = \rho_1(\mathbf{r}, \mathbf{r}) = 2 \sum_{w \in \{\chi_o\}} \chi_w(\mathbf{r}) \chi_w^*(\mathbf{r}), \quad \rho_1 \longrightarrow \rho_s, \quad (16)$$

and we use Eq. (9); the right side of this equation indicates that ρ_s is determined by ρ_1 .

The non-local operator $\hat{v}_{xc}$ can be identified as the exchange-correlation operator. If $|\rho_1\rangle$ is the Kohn-Sham noninteracting state, this operator is local and it implicitly depends only on

ρ_s , the density from $|\rho_1\rangle$ and the interacting state $|\Psi_{Nv}\rangle$. For Brueckner orbital theory,^{40,45} this operator is nonlocal and may depend on the external potential v . For Hartree–Fock theory, the correlation portion of $\hat{v}_{xc}$ is absent; so, in that case, $\hat{v}_{xc}^{\rho_1}$ is just the exchange operator $\hat{K}$, where the kernel of this operator is

$$K_{\rho_1}(\mathbf{r}_1, \mathbf{r}_2) = -\frac{1}{2}r_{12}^{-1}\rho_1(\mathbf{r}_1, \mathbf{r}_2). \quad (17)$$

For latter use, we also write down the expression for the Fock operator:

$$\hat{F}_{\rho_1} = -\frac{1}{2}\nabla^2 + v + \hat{J}_{\rho_s} + \hat{K}_{\rho_1}. \quad (18)$$

III. THE INTERACTING STATE

Now consider an interacting system of electrons. The electronic energy, say $\mathcal{E}_{Nv}$, for an eigenstate of an interacting Hamiltonian is determined by the number of electrons N and the one-body external potential v . This must be the case, since the non-relativistic, electronic Hamiltonian is entirely specified by N and v :

$$\hat{H}_{Nv} = \sum_i^N -\frac{1}{2}\nabla_i^2 + \sum_i^N v(i) + \frac{1}{2} \sum_{i \neq j}^N \sum_j^N r_{ij}^{-1}, \quad (19)$$

where r_{ij}^{-1} is the electron-electron interaction operator. Furthermore, since each N and v yields a different ground state wave function $|\Psi_{Nv}\rangle$.¹ Therefore, the ground state energies $\mathcal{E}_{Nv}$, given by

$$\mathcal{E}_{Nv} = \frac{\langle \Psi_{Nv} | \hat{H}_{Nv} | \Psi_{Nv} \rangle}{\langle \Psi_{Nv} | \Psi_{Nv} \rangle}, \quad (20)$$

are unique functionals of N and v .

Using a second quantization approach, the Hamiltonian does not depend on N , and it can be expressed by

$$\hat{H}_v = \sum_{ij} (i | (-\frac{1}{2}\nabla^2) | j) \hat{E}_{ij} + \sum_{ij} (i | v | j) \hat{E}_{ij} + \frac{1}{2} \sum_{ijkl} (ij | kl) \hat{E}_{ijkl}, \quad (21)$$

where the symmetry-adapted excitation operators are given by Eq. (6) and

$$\hat{E}_{ijkl} = \sum_{\sigma\lambda} a_{i\sigma}^\dagger a_{k\lambda}^\dagger a_{l\lambda} a_{j\sigma}, \quad (22)$$

and the two electrons integrals are spin-free and are written in chemist's notation.¹⁸

IV. THE CORRELATION ENERGY AS IN EXPLICIT FUNCTIONAL OF v , ρ_1 AND v_s

Using many body perturbation theory,^{16,17,18,35} we partition our Hamiltonian into the zeroth order part and a perturbation V :

$$\hat{H}_v = \hat{H}_{v_s} + V_{\rho_1}, \quad (23)$$

where we choose $\hat{H}_{v_s}$, from Eq. (4), as our zeroth-order Hamiltonian. Using Eqs. (2), (4), (21), and (23), we have

$$V_{\rho_1} = \frac{1}{2} \sum_{ijkl} (ij|kl) \hat{E}_{ijkl} + \sum_{ij} (i|v|j) \hat{E}_{ij} - \sum_{ij} (i|v_s|j) \hat{E}_{ij}. \quad (24)$$

The energy through the first order—the zeroth-order energy plus the correction from the 1st order—for many body perturbation theory is independent of the zeroth-order Hamiltonian. This energy functional, given by

$$E_1[\rho_1, v] = \langle \rho_1 | H_v | \rho_1 \rangle, \quad (25)$$

depends on the two functions: the external potential v and the one-particle density matrix ρ_1 . Explicitly, this functional is given by

$$\begin{aligned} E_1[\rho_1, v] = & \int d\mathbf{r}_1 \left[-\frac{1}{2} \nabla_1^2 \rho_1(\mathbf{r}_1, \mathbf{r}_2) \right]_{\mathbf{r}_2=\mathbf{r}_1} \\ & + \int d\mathbf{r}_1 v(\mathbf{r}_1) \rho(\mathbf{r}_1) + \frac{1}{2} \int \int r_{12}^{-1} d\mathbf{r}_1 d\mathbf{r}_2 \rho(\mathbf{r}_1) \rho(\mathbf{r}_2) - \frac{1}{4} \int \int r_{12}^{-1} d\mathbf{r}_1 d\mathbf{r}_2 \rho_1(\mathbf{r}_1, \mathbf{r}_2) \rho_1(\mathbf{r}_2, \mathbf{r}_1). \end{aligned} \quad (26)$$

In many-body perturbation theory, the electronic energy is given by an order-by-order expansion

$$\mathcal{E} = \mathcal{E}_0 + \mathcal{E}_1 + \mathcal{E}_2 + \mathcal{E}_3 + \cdots, \quad (27)$$

where the energy through the first order is

$$E_1 = \mathcal{E}_0 + \mathcal{E}_1, \quad (28)$$

and the correlation energy $\mathcal{E}_{\text{co}}$ includes everything beyond the first order:

$$\mathcal{E}_{\text{co}} = \mathcal{E} - E_1 = \mathcal{E}_2 + \mathcal{E}_3 + \cdots. \quad (29)$$

In order to obtain the an order-by-order expansion of the electronic energy $\mathcal{E}$, it is convenient to write the perturbation V_{ρ_1} , given by Eq. (24), in normal ordered form^{16,17} using $|\rho_1\rangle$ as the vacuum state:

$$V_{\rho_1} = V_c^{\rho_1} + V_1^{\rho_1} + V_2^{\rho_1}, \quad (30)$$

where the constant, one-, and two-body terms are given by

$$V_c^{\rho_1} = \mathcal{E}_1 = E_1 - \mathcal{E}_0, \quad (31)$$

$$V_1^{\rho_1} = - \sum_{ij} (i|\hat{v}_{\text{co}}^{\rho_1}|j) \{\hat{E}_{ij}\}_{\rho_1}, \quad (32)$$

$$V_2^{\rho_1} = \frac{1}{2} \sum_{ijkl} (ij|kl) \{\hat{E}_{ijkl}\}_{\rho_1}, \quad (33)$$

where the correlation potential is defined by Eq. (14), i.e.,

$$\hat{v}_{\text{co}}^{\rho_1} = v_s - v - \hat{J}_{\rho_s} - \hat{K}_{\rho_1}, \quad (34)$$

and $\mathcal{E}_0$, E_1 , $\hat{J}$, and $\hat{K}$ are given by Eqs. (11), (26), (15), and (17), respectively; the brackets $\{\cdots\}_{\rho_1}$ indicate that the excitation operators are in normal ordered form with respect to the vacuum state $|\rho_1\rangle$. Furthermore, subtracting Eq. (18) from (2) and using Eq. (34), we have

$$\hat{v}_{\text{co}}^{\rho_1} = \hat{\mathcal{F}}_{\rho_1} - \hat{F}_{\rho_1}. \quad (35)$$

Eq. (32) is the most convenient expression for V_1 . However, we can also substitute Eq. (34) into Eq. (32), yielding individual diagrams from the perturbation expansion that depend explicitly on v and v_s . Substituting Eq. (35) into Eq. (32) separates V_1 into the zeroth-order part $\hat{\mathcal{F}}$ and the Fock-operator part $\hat{F}$. (Note that $\{\hat{F}\}_{\rho_1}$ is the one-body portion of the Hamiltonian.)⁴¹ While the individual terms from the perturbation expansion can depend on v_s or $\hat{\mathcal{F}}$, the correlation energy $\mathcal{E}_{\text{co}}$, unless approximated from a truncated expansion, cannot depend on either v_s or $\hat{\mathcal{F}}$; we now prove that $\mathcal{E}_{\text{co}}$ is an implicit functional of v and ρ_1 .

Consider the electronic energy $\mathcal{E}$, determined by v and N , while E_1 is an explicit functional of v and ρ_1 . Hence, the correlation energy, given by $(\mathcal{E}_{\text{co}} = \mathcal{E} - E_1)$, is determined by v , ρ_1 , and N , or just v and ρ_1 , since ρ_1 determines N . Therefore, as mentioned above, at least for the ground state, the correlation energy is an implicit functional of v and ρ_1 , i.e., $\mathcal{E}_{\text{co}}[v, \rho_1]$. We now demonstrate that the individual perturbation terms, or diagrams, can be written as explicit functional of v , v_s and ρ_1 .




FIG. 1: The three diagrams from the second-order correlation-energy $\mathcal{E}_{\text{co}}$ and two other diagrams.

Generally speaking, perturbation theory expresses $\mathcal{E}_{\text{co}}$ as a functional of the occupied and unoccupied orbitals from $|\rho_1\rangle$, their orbitals energies $\{\epsilon\}$, and the one-body portion or the perturbation V_1 , which, in our case is $-\hat{v}_{\text{co}}^{\rho_1}$, $(v - v_s + \hat{J}_{\rho_s} + \hat{K}_{\rho_1})$, or $(\hat{F} - \hat{\mathcal{F}})$, as indicated in Eqs. (32), (34), and (35). The correlation-energy diagrams, however, do not depend on the constant portion of the perturbation V_c ; also, the two-body part V_2 is universal, since this operator is the same for all electronic systems.

Consider for the second-order (correction to the) energy, given by

$$\mathcal{E}_2 = \mathcal{E}_{2u} + \mathcal{E}_{2d} + \mathcal{E}_{2x}, \quad (36)$$

where the Goldstone diagrams for these terms are given by the first three entries of Fig. 1, respectively. For the first term, we have

$$\mathcal{E}_{2u}[\chi_o, \chi_u, \epsilon, v_{\text{co}}^{\rho_1}] = 2 \sum_{w \in \{\chi_o\}} \sum_{r \in \{\chi_u\}} \frac{1}{\epsilon_w - \epsilon_r} (r | \hat{v}_{\text{co}}^{\rho_1} | w) (w | \hat{v}_{\text{co}}^{\rho_1} | r), \quad (37)$$

where this diagram depends explicitly on $v_{\text{co}}^{\rho_1}$ —the kernel of $\hat{v}_{\text{co}}^{\rho_1}$ —the orbital energies $\{\epsilon\}$, and both the occupied and unoccupied orbitals, $\{\chi_o\}$ and $\{\chi_u\}$. Writing out the integrations explicitly, we have

$$\mathcal{E}_{2u}[\chi_o, \chi_u, \epsilon, v_{\text{co}}^{\rho_1}] = 2 \sum_{wr} \int \int d\mathbf{r}_1 d\mathbf{r}_2 \frac{1}{\epsilon_w - \epsilon_r} \hat{v}_{\text{co}}^{\rho_1}(\mathbf{r}_1) \chi_w(\mathbf{r}_1) \chi_w^*(\mathbf{r}_2) \hat{v}_{\text{co}}^{\rho_1}(\mathbf{r}_2) \chi_r(\mathbf{r}_2) \chi_r^*(\mathbf{r}_1). \quad (38)$$

In order to remove the energy denominator, we use the following Laplace transform:^{19,27}

$$\frac{1}{\epsilon_w - \epsilon_r} = \int_0^\infty dt e^{-t(\epsilon_w - \epsilon_r)}, \quad (39)$$

giving

$$\begin{aligned} \mathcal{E}_{2u}[\chi_o, \chi_u, \epsilon, \hat{v}_{\text{co}}^{\rho_1}] = \\ 2 \sum_{wr} \int \int d\mathbf{r}_1 d\mathbf{r}_2 \int_0^\infty dt_1 e^{t_1 \epsilon_r} \chi_r^*(\mathbf{r}_1) \hat{v}_{\text{co}}^{\rho_1}(\mathbf{r}_1) e^{-t_1 \epsilon_w} \chi_w(\mathbf{r}_1) \chi_w^*(\mathbf{r}_2) \hat{v}_{\text{co}}^{\rho_1}(\mathbf{r}_2) \chi_r(\mathbf{r}_2). \end{aligned} \quad (40)$$

Using Eqs. (12) and (13), and the observation that $\hat{\mathcal{F}}_{\hat{v}_s}$ depends explicitly on the kernel v_s , and, according to Eq. (34), $\hat{v}_{\text{co}}^{\rho_1}$ depends explicitly on v and v_s , we have

$$\mathcal{E}_{2u}[\chi_o, \chi_u, v, v_s] = \quad (41)$$

$$2 \sum_{wr} \int \int d\mathbf{r}_1 d\mathbf{r}_2 \int_0^\infty dt_1 \left(e^{t_1 \hat{\mathcal{F}}_{v_s}^1} \chi_r^*(\mathbf{r}_1) \right) \hat{v}_{\text{co}}^{\rho_1}(\mathbf{r}_1) \left(e^{-t_1 \hat{\mathcal{F}}_{v_s}^1} \chi_w(\mathbf{r}_1) \right) \chi_w^*(\mathbf{r}_2) \hat{v}_{\text{co}}^{\rho_1}(\mathbf{r}_2) \chi_r(\mathbf{r}_2),$$

where $\hat{\mathcal{F}}_{v_s}^i$ denotes $\hat{\mathcal{F}}_{v_s}(\mathbf{r}_i)$ and the round brackets $(\dots)$ serve to remind us that $e^{t_1 \hat{\mathcal{F}}_{v_s}^1}$ acts exclusively upon $\chi_r^*(\mathbf{r}_1)$ and $e^{-t_1 \hat{\mathcal{F}}_{v_s}^1}$ acts upon $\chi_w(\mathbf{r}_1)$. Note that, since $\hat{v}_s$ and $-\frac{1}{2}\nabla^2$ do not commute, the exponential operator, defined by

$$e^{t \hat{\mathcal{F}}_{v_s}} = \exp \left[t \left(-\frac{1}{2}\nabla^2 + \hat{v}_s \right) \right] = 1 + t \left(-\frac{1}{2}\nabla^2 + \hat{v}_s \right) + \frac{t^2}{2!} \left(-\frac{1}{2}\nabla^2 + \hat{v}_s \right) \left(-\frac{1}{2}\nabla^2 + \hat{v}_s \right) + \dots, \quad (42)$$

cannot separate into two factors:⁴⁶

$$\exp \left[t \left(-\frac{1}{2}\nabla^2 + \hat{v}_s \right) \right] \neq \exp \left[t \left(-\frac{1}{2}\nabla^2 \right) \right] \times \exp \left[t \hat{v}_s \right], \quad \text{if } \left[-\frac{1}{2}\nabla^2, \hat{v}_s \right] \neq 0. \quad (43)$$

In order to make our formulas less cluttered, especially when considering high order diagrams, we introduce a notation where the brackets are understood:

$$\mathcal{E}_{2u}[\chi_o, \chi_u, v, v_s] = \quad (44)$$

$$2 \sum_{wr} \int \int d\mathbf{r}_1 d\mathbf{r}_2 \int_0^\infty dt_1 e^{t_1 \tilde{\mathcal{F}}_{v_s}^{1*}} \chi_r^*(\mathbf{r}_1) \hat{v}_{\text{co}}^{\rho_1}(\mathbf{r}_1) e^{-t_1 \tilde{\mathcal{F}}_{v_s}^1} \chi_w(\mathbf{r}_1) \chi_w^*(\mathbf{r}_2) \hat{v}_{\text{co}}^{\rho_1}(\mathbf{r}_2) \chi_r(\mathbf{r}_2),$$

where $\tilde{\mathcal{F}}_{v_s}$ is identical to $\hat{\mathcal{F}}_{v_s}$, except that it acts exclusively on χ ; it does not act on the correlation operator $\hat{v}_{\text{co}}^{\rho_1}$ (or its kernel $v_{\text{co}}^{\rho_1}$) or the complex conjugate function χ^* . Similarly, $\tilde{\mathcal{F}}_{v_s}^*$ is identical to $\hat{\mathcal{F}}_{v_s}$, except that it acts exclusively on χ^* ; it does not act on $v_{\text{co}}^{\rho_1}$ or χ . The following identities clarify these definitions:

$$e^{-t_1 \tilde{\mathcal{F}}_{v_s}^1} \chi^*(\mathbf{r}_1) \hat{v}_{\text{co}}^{\rho_1}(\mathbf{r}_1) \chi(\mathbf{r}_1) = \chi^*(\mathbf{r}_1) \hat{v}_{\text{co}}^{\rho_1}(\mathbf{r}_1) e^{-t_1 \hat{\mathcal{F}}_{v_s}^1} \chi(\mathbf{r}_1), \quad (45)$$

$$e^{-t_1 \tilde{\mathcal{F}}_{v_s}^{1*}} \chi(\mathbf{r}_1) \hat{v}_{\text{co}}^{\rho_1}(\mathbf{r}_1) \chi^*(\mathbf{r}_1) = \chi(\mathbf{r}_1) \hat{v}_{\text{co}}^{\rho_1}(\mathbf{r}_1) e^{-t_1 \hat{\mathcal{F}}_{v_s}^1} \chi^*(\mathbf{r}_1). \quad (46)$$

In order to further compress our notation, we define the following transformed correlation operator:

$$\tilde{v}_{\text{co}\rho_1}^{v_s \mathbf{z}_i} = e^{t_i \tilde{\mathcal{F}}_{v_s}^{i*}} \hat{v}_{\text{co}}^{\rho_1}(\mathbf{r}_i) e^{-t_i \tilde{\mathcal{F}}_{v_s}^i}, \quad (47)$$

where $\mathbf{z}_i$ is a combined index of $\mathbf{r}_i$ and t_i , i.e., $\mathbf{z}_i = \mathbf{r}_i, t_i$. This notation yields

$$\mathcal{E}_{2u}[\rho_1, \kappa_{\rho_1}, v, v_s] = \frac{1}{2} \int \int d\mathbf{r}_1 d\mathbf{r}_2 \int_0^\infty dt_1 \tilde{v}_{\text{co}\rho_1}^{v_s \mathbf{z}_1} \rho_1(\mathbf{r}_1, \mathbf{r}_2) \hat{v}_{\text{co}}^{\rho_1}(\mathbf{r}_2) \kappa_{\rho_1}(\mathbf{r}_2, \mathbf{r}_1), \quad (48)$$

where ρ_1 is given by Eq. (9), while its orthogonal complement, κ_{ρ_1} , is given by

$$\kappa_{\rho_1}(\mathbf{r}_1, \mathbf{r}_2) = 2 \sum_{r \in \{\chi_u\}} \chi_r(\mathbf{r}_1) \chi_r^*(\mathbf{r}_2), \quad (49)$$

and the sum is over the unoccupied orbitals $\{\chi_u\}$. For a complete set of basis functions, we have

$$\delta(\mathbf{r}_1 - \mathbf{r}_2) = \sum_i \chi_i(\mathbf{r}_1) \chi_i^*(\mathbf{r}_2), \quad i \in \{\chi_o\} \cup \{\chi_u\}, \quad (50)$$

and therefore

$$\delta(\mathbf{r}_1 - \mathbf{r}_2) = \frac{1}{2} (\rho_1(\mathbf{r}_1, \mathbf{r}_2) + \kappa_{\rho_1}(\mathbf{r}_1, \mathbf{r}_2)). \quad (51)$$

So we can remove κ_{ρ_1} from Eq. (48), giving

$$\begin{aligned} \mathcal{E}_{2u}[\rho_1, v, v_s] = & \int \int d\mathbf{r}_1 d\mathbf{r}_2 \int_0^\infty dt_1 \tilde{v}_{\text{co}\rho_1}^{v_s \mathbf{z}_1} \rho_1(\mathbf{r}_1, \mathbf{r}_2) \hat{v}_{\text{co}}^{\rho_1}(\mathbf{r}_2) \delta(\mathbf{r}_2 - \mathbf{r}_1) \\ & - \frac{1}{2} \int \int d\mathbf{r}_1 d\mathbf{r}_2 \int_0^\infty dt_1 \tilde{v}_{\text{co}\rho_1}^{v_s \mathbf{z}_1} \rho_1(\mathbf{r}_1, \mathbf{r}_2) \hat{v}_{\text{co}}^{\rho_1}(\mathbf{r}_2) \rho_1(\mathbf{r}_2, \mathbf{r}_1). \end{aligned} \quad (52)$$

Hence, we have $\mathcal{E}_{2u}$ as an explicit functional v , v_s , and ρ_1 .

In order to evaluate Eq. (52), one must handle the operator $\exp(t\hat{\mathcal{F}})$ acting on the Dirac delta function δ , since the first term on the right side is given by

$$\begin{aligned} & \int \int d\mathbf{r}_1 d\mathbf{r}_2 \int_0^\infty dt_1 \tilde{v}_{\text{co}\rho_1}^{v_s \mathbf{z}_1} \rho_1(\mathbf{r}_1, \mathbf{r}_2) \hat{v}_{\text{co}}^{\rho_1}(\mathbf{r}_2) \delta(\mathbf{r}_2 - \mathbf{r}_1) \\ & = \int \int d\mathbf{r}_1 d\mathbf{r}_2 \int_0^\infty dt_1 \hat{v}_{\text{co}}^{\rho_1}(\mathbf{r}_1) \left(e^{-t_1 \hat{\mathcal{F}}_{v_s}^1} \rho_1(\mathbf{r}_1, \mathbf{r}_2) \right) \hat{v}_{\text{co}}^{\rho_1}(\mathbf{r}_2) \left(e^{t_1 \hat{\mathcal{F}}_{v_s}^1} \delta(\mathbf{r}_2 - \mathbf{r}_1) \right). \end{aligned} \quad (53)$$

Using the expansion for $e^{t\hat{\mathcal{F}}}$ given by Eq. (42), this factor will yield terms containing the Laplacian operator ∇^2 acting on δ , i.e., $\nabla^2 \delta$, as well as other higher order terms. e.g, $\nabla^4 \delta$. One way to handle these terms is by using a basis set expansion for δ , as in Eq. (50), where any complete basis can be used that satisfy the boundary conditions. Another possibility is to use one of the integral representations of the delta function, for example,

$$\delta(\mathbf{r}_1 - \mathbf{r}_2) = \frac{1}{8\pi^3} \int e^{i\mathbf{k} \cdot (\mathbf{r}_1 - \mathbf{r}_2)} d\mathbf{k}. \quad (54)$$

Now consider the second term of Eq. (36), given by the second diagram from Fig. 1; evaluating this diagram yields the following expression:

$$\mathcal{E}_{2d}[\chi_o, \chi_u, \epsilon] = 2 \sum_{w \in \{\chi_o\}} \sum_{x \in \{\chi_o\}} \sum_{r \in \{\chi_u\}} \sum_{s \in \{\chi_u\}} \frac{1}{\epsilon_w - \epsilon_r + \epsilon_x - \epsilon_s} (rw|sx)(wr|xs), \quad (55)$$

and this term can be written as

$$\begin{aligned} \mathcal{E}_{2d}[\chi_o, \chi_u, \epsilon] &= 2 \sum_{wx} \sum_{rs} (\epsilon_w - \epsilon_r + \epsilon_x - \epsilon_s)^{-1} \\ &\times \int \int \int \int d\mathbf{r}_1 d\mathbf{r}_2 d\mathbf{r}_3 d\mathbf{r}_4 r_{12}^{-1} \chi_r^*(\mathbf{r}_1) \chi_w(\mathbf{r}_1) \chi_s^*(\mathbf{r}_2) \chi_x(\mathbf{r}_2) \chi_w^*(\mathbf{r}_3) \chi_r(\mathbf{r}_3) r_{34}^{-1} \chi_x^*(\mathbf{r}_4) \chi_s(\mathbf{r}_4). \end{aligned} \quad (56)$$

Using a generalization of Eq. (39), we have

$$\begin{aligned} \mathcal{E}_{2d}[\chi_o, \chi_u, \epsilon] &= 2 \sum_{wx} \sum_{rs} \int \int \int \int d\mathbf{r}_1 d\mathbf{r}_2 d\mathbf{r}_3 d\mathbf{r}_4 \int_0^\infty dt_{12} \\ &\times e^{t_{12}\epsilon_r} e^{t_{12}\epsilon_s} r_{12}^{-1} e^{-t_{12}\epsilon_w} e^{-t_{12}\epsilon_x} \chi_r^*(\mathbf{r}_1) \chi_w(\mathbf{r}_1) \chi_s^*(\mathbf{r}_2) \chi_x(\mathbf{r}_2) \chi_w^*(\mathbf{r}_3) \chi_r(\mathbf{r}_3) r_{34}^{-1} \chi_x^*(\mathbf{r}_4) \chi_s(\mathbf{r}_4). \end{aligned} \quad (57)$$

Using our notation, defined by Eqs. (45) and (46), and Eqs. (49) and (9), we obtain

$$\begin{aligned} \mathcal{E}_{2d}[\rho_1, v_s] &= \frac{1}{8} \int \int \int \int d\mathbf{r}_1 d\mathbf{r}_2 d\mathbf{r}_3 d\mathbf{r}_4 \int_0^\infty dt_{12} \\ &\times (\tilde{r}_{12}^{\hat{v}_s})^{-1} \rho_1(\mathbf{r}_1, \mathbf{r}_3) \kappa_{\rho_1}(\mathbf{r}_3, \mathbf{r}_1) \rho_1(\mathbf{r}_2, \mathbf{r}_4) \kappa_{\rho_1}(\mathbf{r}_4, \mathbf{r}_2) r_{34}^{-1}, \end{aligned} \quad (58)$$

where the transformed electron-electron interaction term $(\tilde{r}_{12}^{\hat{v}_s})^{-1}$ depends on the variables t_{12} , $\mathbf{r}_1$, $\mathbf{r}_2$, and the function v_s ; this operator has the following form:

$$\frac{1}{\tilde{r}_{12}^{\hat{v}_s}} = e^{t_{12}\tilde{\mathcal{F}}_{v_s}^{1*}} e^{t_{12}\tilde{\mathcal{F}}_{v_s}^{2*}} r_{12}^{-1} e^{-t_{12}\tilde{\mathcal{F}}_{v_s}^1} e^{-t_{12}\tilde{\mathcal{F}}_{v_s}^2}. \quad (59)$$

The $2d$ subscript indicates that $\mathcal{E}_{2d}$ correspond to a second order “direct” diagram. The second-order exchange term $\mathcal{E}_{2x}$, from Eq. (36), is represented by the third diagram of Fig. 1. This diagram can be evaluated in a similar way as the direct one $\mathcal{E}_{2d}$, giving

$$\begin{aligned} \mathcal{E}_{2x}[\rho_1, v_s] &= -\frac{1}{16} \int \int \int \int d\mathbf{r}_1 d\mathbf{r}_2 d\mathbf{r}_3 d\mathbf{r}_4 \int_0^\infty dt_{12} \\ &(\tilde{r}_{12}^{\hat{v}_s})^{-1} \rho_1(\mathbf{r}_1, \mathbf{r}_3) \kappa_{\rho_1}(\mathbf{r}_3, \mathbf{r}_2) \rho_1(\mathbf{r}_2, \mathbf{r}_4) \kappa_{\rho_1}(\mathbf{r}_4, \mathbf{r}_1) r_{34}^{-1}. \end{aligned} \quad (60)$$

Note that the $\mathcal{E}_{2d}$ and $\mathcal{E}_{2x}$ do not explicitly depend on the external potential v .

In Fig. 1, the dummy indices that are used are denoted in the diagrams. For example, the first diagram has $\mathbf{z}_1$ for the first vertex and $\mathbf{r}_2$ for the second index, where these choices are indicated by 1 and 2 in the figure. In order to evaluate this diagram and obtain Eq. (48) directly, we start at the bottom vertex and go backwards: The bottom vertex gives a factor of $-\tilde{v}_{\text{co}\rho_1}^{v_s \mathbf{z}_1}$; the hole line from vertex 1 to 2 gives a factor of $\frac{1}{2}\rho_1(\mathbf{r}_1, \mathbf{r}_2)$; the top vertex gives

$-\hat{v}_{co}^{\rho_1}(\mathbf{r}_2)$; the particle line from vertex 2 to 1 gives $\frac{1}{2}\kappa_{\rho_1}(\mathbf{r}_2, \mathbf{r}_1)$. Furthermore, there is an additional factor of two for the loop, since the spin state is conserved on an orientated path.¹⁷ Similarly, the second two diagrams from the figure can be evaluated to give Eqs. (58) and (60), where in both diagrams, the lower interactions, with dummy indices $\mathbf{r}_1$, $\mathbf{r}_2$ and t_{12} , yield the operator $(\tilde{r}_{12}^{\hat{v}_s})^{-1}$. For the second diagram in the figure, the direct one $\mathcal{E}_{2d}$, we have the following factors: a $\frac{1}{2}$, since the diagram is symmetric, a factor of 2×2 for the two loops, and a factor of $\frac{1}{2} \times \frac{1}{2} \times \frac{1}{2} \times \frac{1}{2}$ for the two ρ_1 terms and two κ terms. Putting this together gives $\frac{1}{2} \times 2 \times 2 \times \frac{1}{2} \times \frac{1}{2} \times \frac{1}{2} \times \frac{1}{2} = \frac{1}{8}$, in agreement with Eq. (58).

Now consider the fourth diagram in Fig. 1, where for convenience we denote this third order diagram by $\mathcal{E}_{3d}^{hh}$, since this is a direct diagram where the middle interaction lies on two hole lines. Evaluating this diagram gives

$$\mathcal{E}_{3d}^{hh} = 2 \sum_{wx} \sum_{rs} \sum_{yz} \frac{(yr|zs)(wy|xz)(rw|sx)}{(\epsilon_y - \epsilon_r + \epsilon_z - \epsilon_s)(\epsilon_w - \epsilon_r + \epsilon_x - \epsilon_s)}, \quad (61)$$

where we have two energy denominators: One corresponding to the lowest interaction, given by $(\epsilon_w + \epsilon_x - \epsilon_r - \epsilon_s)$, and one for the middle interaction, given by $(\epsilon_y + \epsilon_z - \epsilon_r - \epsilon_s)$, where we denote the occupied orbitals by w , x , y , and z and the unoccupied orbitals by r and s . Using the following identity:¹⁹

$$\frac{1}{\epsilon_j \epsilon_i} = \int_0^\infty dt_2 e^{-t_2(\epsilon_j - \epsilon_i)} \int_{t_2}^\infty dt_1 e^{-t_1 \epsilon_i}, \quad (62)$$

the product of the two energy denominator can be written as

$$\begin{aligned} & \frac{1}{(\epsilon_y - \epsilon_r + \epsilon_z - \epsilon_s)(\epsilon_w - \epsilon_r + \epsilon_x - \epsilon_s)} \\ &= \int_0^\infty dt_{34} e^{-t_{34}(\epsilon_y - \epsilon_w + \epsilon_z - \epsilon_x)} \int_{t_{34}}^\infty dt_{12} e^{-t_{12}(\epsilon_w - \epsilon_r + \epsilon_x - \epsilon_s)}. \end{aligned} \quad (63)$$

Comparing the right side of this expression with the the fourth diagram in the figure, or with Eq. (61), we see that the two lowest interactions, with the following two matrix elements: $(rw|sx)$ and $(wy|xz)$, are directly related to the energy exponential factors: $t_{12}(\epsilon_r - \epsilon_w + \epsilon_s - \epsilon_x)$ and $t_{34}(\epsilon_w - \epsilon_y + \epsilon_x - \epsilon_z)$. Therefore, multiplying Eq. (63) by these two matrix

elements gives

$$\begin{aligned}
& \frac{(rw|sx)(wy|xz)}{(\epsilon_y - \epsilon_r + \epsilon_z - \epsilon_s)(\epsilon_w - \epsilon_r + \epsilon_x - \epsilon_s)} = \int \int \int \int d\mathbf{r}_1 d\mathbf{r}_2 d\mathbf{r}_3 d\mathbf{r}_4 \\
& \quad \times \int_0^\infty dt_{34} \int_{t_{34}}^\infty dt_{12} \left(e^{t_{12}\epsilon_r} e^{t_{12}\epsilon_s} r_{12}^{-1} e^{-t_{12}\epsilon_w} e^{-t_{12}\epsilon_x} \chi_r^*(\mathbf{r}_1) \chi_w(\mathbf{r}_1) \chi_s^*(\mathbf{r}_2) \chi_x(\mathbf{r}_2) \right) \\
& \quad \times \left(e^{t_{34}\epsilon_w} e^{t_{34}\epsilon_x} r_{34}^{-1} e^{-t_{34}\epsilon_y} e^{-t_{34}\epsilon_z} \chi_w^*(\mathbf{r}_3) \chi_y(\mathbf{r}_3) \chi_x^*(\mathbf{r}_4) \chi_z(\mathbf{r}_4) \right) \\
& = \int \int \int \int d\mathbf{r}_1 d\mathbf{r}_2 d\mathbf{r}_3 d\mathbf{r}_4 \int_0^\infty dt_{34} \int_{t_{34}}^\infty dt_{12} \\
& \quad \times (\tilde{r}_{12}^{\hat{v}_s})^{-1} \chi_r^*(\mathbf{r}_1) \chi_w(\mathbf{r}_1) \chi_s^*(\mathbf{r}_2) \chi_x(\mathbf{r}_2) (\tilde{r}_{34}^{\hat{v}_s})^{-1} \chi_w^*(\mathbf{r}_3) \chi_y(\mathbf{r}_3) \chi_x^*(\mathbf{r}_4) \chi_z(\mathbf{r}_4),
\end{aligned} \tag{64}$$

where we used Eq. (59). Comparing Eqs. (64) and (61), and using Eqs. (9) and (49), gives

$$\begin{aligned}
\mathcal{E}_{3d}^{\text{hh}}[\rho_1, v_s] &= \frac{1}{32} \int \int \int \int \int \int d\mathbf{r}_1 d\mathbf{r}_2 d\mathbf{r}_3 d\mathbf{r}_4 d\mathbf{r}_5 d\mathbf{r}_6 \int_0^\infty dt_{34} \int_{t_{34}}^\infty dt_{12} \\
& \quad \times (\tilde{r}_{12}^{\hat{v}_s})^{-1} (\tilde{r}_{34}^{\hat{v}_s})^{-1} r_{56}^{-1} \rho_1(\mathbf{r}_1, \mathbf{r}_3) \rho_1(\mathbf{r}_3, \mathbf{r}_5) \kappa_{\rho_1}(\mathbf{r}_5, \mathbf{r}_1) \rho_1(\mathbf{r}_2, \mathbf{r}_4) \rho_1(\mathbf{r}_4, \mathbf{r}_6) \kappa_{\rho_1}(\mathbf{r}_6, \mathbf{r}_2).
\end{aligned} \tag{65}$$

Note that in this diagram the two bottom interactions, r_{12}^{-1} and r_{34}^{-1} , are transformed into $(\tilde{r}_{12}^{\hat{v}_s})^{-1}$ and $(\tilde{r}_{34}^{\hat{v}_s})^{-1}$, respectively, while the top interaction, r_{56}^{-1} , is not transformed. Furthermore, the transformed interaction that is next to the top, $(\tilde{r}_{34}^{\hat{v}_s})^{-1}$, with the dummy index t_{34} , gives an integration limit of 0 to ∞ , while the one below that, $(\tilde{r}_{12}^{\hat{v}_s})^{-1}$, with the dummy index t_{12} , has an integration limit of t_{34} to ∞ . By considering higher-order diagrams, and by generalizing the identity given by Eq. (62), e.g.,¹⁹

$$\frac{1}{\varepsilon_3 \varepsilon_2 \varepsilon_1} = \int_0^\infty dt_3 e^{-t_2(\varepsilon_3 - \varepsilon_2)} \int_{t_3}^\infty dt_2 e^{-t_2(\varepsilon_2 - \varepsilon_1)} \int_{t_2}^\infty dt_1 e^{-t_1 \varepsilon_1}, \tag{66}$$

it is easily shown that the following rules apply to all diagram: 1) A factor of $\frac{1}{2}\rho_1$ and $\frac{1}{2}\kappa_{\rho_1}$ is given for each hole and particle line, respectively. 2) Interactions given by $\hat{v}_{\text{co}}^{\rho_1}(\mathbf{r}_i)$ and r_{ij}^{-1} are transformed into $\tilde{v}_{\text{co}\rho_1}^{v_s \mathbf{z}_i}$ and $(\tilde{r}_{ij}^{\hat{v}_s})^{-1}$, unless they are the last, or uppermost, interaction. 3) The spatial dummy indices, $\mathbf{r}_1, \mathbf{r}_2, \dots$, are integrated over. 4) A t_i dummy index from $\tilde{v}_{\text{co}\rho_1}^{v_s \mathbf{z}_i}$ and a t_{ij} index from $(\tilde{r}_{ij}^{\hat{v}_s})^{-1}$ are integrated over using the following rules to obtain their integration limits: i) If the interaction—either $\tilde{v}_{\text{co}\rho_1}^{v_s \mathbf{z}_i}$ or $(\tilde{r}_{ij}^{\hat{v}_s})^{-1}$ —is the second from the top—the next to the last one—the integration limit is from 0 to ∞ . ii) For other interactions, the integration limit is from t_j to ∞ , if the interaction above it is $\tilde{v}_{\text{co}\rho_1}^{v_s \mathbf{z}_j}$; while, the integration limit is from t_{jk} to ∞ , if the interaction above it is $(\tilde{r}_{jk}^{\hat{v}_s})^{-1}$. In addition, we also have the usual rules of symmetry and sign factors as well as a factor of two for each loop¹⁷ corresponding to a sum

over each spin state, α and β . Also, according to Eq. (32), the one-body interaction has a negative sign associated with it, it is $-\tilde{v}_{\text{co}\rho_1}^{v_s \mathbf{z}_1}$.

As a final example, consider the last diagram appearing in Fig. 1, where we denote this arbitrary (arb) fourth order diagram by $\mathcal{E}_{\text{arb}}$. Using the rules above we get

$$\mathcal{E}_{\text{arb}}[\rho_1, v, v_s] = -\frac{1}{32} \int \int \int \int \int \int \int d\mathbf{r}_1 d\mathbf{r}_2 d\mathbf{r}_3 d\mathbf{r}_4 d\mathbf{r}_5 d\mathbf{r}_6 d\mathbf{r}_7 \int_0^\infty dt_5 \int_{t_5}^\infty dt_{34} \int_{t_{34}}^\infty dt_{12} \quad (67)$$

$$\times (\tilde{r}_{12}^{\hat{v}_s})^{-1} (\tilde{r}_{34}^{\hat{v}_s})^{-1} \tilde{v}_{\text{co}\rho_1}^{v_s \mathbf{z}_5} r_{67}^{-1} \rho_1(\mathbf{r}_1, \mathbf{r}_3) \rho_1(\mathbf{r}_3, \mathbf{r}_6) \kappa_{\rho_1}(\mathbf{r}_6, \mathbf{r}_1) \rho_1(\mathbf{r}_2, \mathbf{r}_4) \rho_1(\mathbf{r}_4, \mathbf{r}_7) \kappa_{\rho_1}(\mathbf{r}_7, \mathbf{r}_5) \kappa_{\rho_1}(\mathbf{r}_5, \mathbf{r}_2).$$

Hence, using the above rules for all diagram, and Eqs. (34) and (51)—for $\hat{v}_{\text{co}}^{\rho_1}$ and κ_{ρ_1} —yields the correlation energy $\mathcal{E}_{\text{co}}$ as an explicit functional ρ_1 , v , and v_s . i.e, $\mathcal{E}_{\text{co}}[\rho_1, v, v_s]$. The other two choices are $\mathcal{E}_{\text{co}}$ as an explicit functional ρ_1 , $v_{\text{co}}^{\rho_1}$ and v_s , or, using Eq. (35), $\mathcal{E}_{\text{co}}$ as an explicit functional ρ_1 , v , and $\hat{\mathcal{F}}_{\rho_1}$. i.e, $\mathcal{E}_{\text{co}}[\rho_1, v, \hat{\mathcal{F}}_{\rho_1}]$, where the v dependence arises from the Fock operator $\hat{F}_{\rho_1}$, using Eq. (18).

It is well known the the correlation energy cannot depend on the zeroth order Hamiltonian $\hat{\mathcal{F}}_{\rho_1}$. Therefore, while the expansion $\mathcal{E}_{\text{co}}[\rho_1, v, \hat{\mathcal{F}}_{\rho_1}]$ has individual diagrams that explicitly depend of $\hat{\mathcal{F}}_{\rho_1}$, the entire series does not. Certain infinite-order partial summations also yield results that are independent of a zeroth order Hamiltonian. For example, if all diagrams are included that correspond to a coupled cluster calculation, the final approximation of $\mathcal{E}_{\text{co}}$ does not depend on the zeroth order Hamiltonian $\hat{\mathcal{F}}_{\rho_1}$, that is, we have the following implicit functional: $\mathcal{E}_{\text{co}}[\rho_1, v]$.

* Electronic address: james.finley@enmu.edu

¹ R. M. Dreizler and E. K. U. Gross, *Density Functional Theory: An Approach to the Quantum Many-Body Problem* (Springer-Verlag, Berlin, Heidelberg, New York, London, Paris, Tokyo, Hong Kong, Barcelona, 1990).

² R. G. Parr and W. Yang, *Density-Functional Theory of Atoms and Molecules* (Oxford University Press, New York, 1989).

³ M. Springborg, ed., *Density Functional Methods in Chemistry and Material Science* (Wiley, Chichester, 1997).

- ⁴ D. E. Ellis, ed., *Density Functional Methods of Molecules, Clusters, and Solids* (Kluwer, Dordrecht, 1995).
- ⁵ E. K. U. Gross and R. M. Dreizler, eds., *Density Functional Theory* (Plenum, New York, 1994).
- ⁶ J. Seminario and P. Politzer, eds., *Modern Density Functional Theory. A Tool for Chemistry* (Elsevier, Amsterdam, 1995).
- ⁷ N. C. Handy, in *Europe Summerschool in Quantum Chemistry*, B. O. Roos and P. -O. Widmark eds. (1997), 2nd ed., (see internet address www.teokem.lu.se/esqc).
- ⁸ P. Hohenberg and W. Kohn, Phys. Rev. **136B**, 864 (1964).
- ⁹ C. Fiolhais, F. Nogueira, and M. Marques, eds., *Lecture Notes in Physics: A primer on Density Functional Theory* (Springer-Verlag, Berlin, Heidelberg, New York, Barcelona, Hong Kong, London, Milan, Paris, Tokyo, 2003), (and references therein).
- ¹⁰ R. T. Sharp and G. K. Horton, Phys. Rev. **90**, 317 (1953).
- ¹¹ J. Talman and W. Shadwick, Phys. Rev. A **14**, 36 (1976).
- ¹² Y. Li, J. B. Krieger, and G. J. Iafrate, Phys. Rev. A **47**, 165 (1993).
- ¹³ V. Shaginyan, Phys. Rev. A **47**, 1507 (1994).
- ¹⁴ A. Görling and M. Levy, Phys. Rev. A **50**, 196 (1994).
- ¹⁵ T. Grabo, T. Kreibich, S. Kurth, and E.K.U Gross, in *Strong Coulomb Correlation in Electronic Structure: Beyond the Local Density Approximation* (Gordon and Breach, Tokyo, 2000).
- ¹⁶ I. Lindgren and J. Morrison, *Atomic Many-Body Theory*, Springer Series on Atoms and Plasmas (Springer-Verlag, New York, Berlin, Heidelberg, 1986), 2nd ed.
- ¹⁷ J. Paldus and J. Čížek, Adv. Quantum Chem. **9**, 105 (1975).
- ¹⁸ A. Szabo and N. S. Ostlund, *Modern Quantum Chemistry: Introduction to Advanced Electronic Structure Theory* (Macmillan, New York, 1982).
- ¹⁹ F. E. Harris, H. J. Monkhorst, and D. L. Freeman, *Algebraic and Diagrammatic Methods in Many-Fermion Theory* (Oxford University Press, New York, 1992).
- ²⁰ P. Pulay and S. Saebø, Theor. Chim. Acta **69**, 357 (1986).
- ²¹ S. Saebø and P. Pulay, J. Chem. Phys. **86**, 914 (1987).
- ²² M. Head-Gordon, P. Maeslen, and C. White, J. Chem. Phys. **108**, 616 (1998).
- ²³ M. Schütz, G. Hetzer, and H.-J. Werner, J. Chem. Phys. **111**, 5691 (1999).
- ²⁴ G. Hetzer, M. Schütz, H. Stoll, and H. -J. Werner, J. Chem. Phys. **113**, 9443 (2000).
- ²⁵ J. P. Finley, Chem. Phys. Lett. **328**, 51 (2000).

- ²⁶ Y. Nakao and K. Hirao, J. Chem. Phys. **120**, 6375 (2004).
- ²⁷ M. Häser and J. Amlöf, J. Chem. Phys. **96**, 489 (1992).
- ²⁸ P. Y. Ayala and G. E. Scuseria, J. Chem. Phys. **111**, 8330 (1999).
- ²⁹ P. Y. Ayala and G. E. Scuseria, J. Chem. Phys. **110**, 3660 (1999).
- ³⁰ A. D. Becke, J. Chem. Phys. **98**, 5648 (1993).
- ³¹ K. Burke, M. Ernzerhof, and J. P. Perdew, Chem. Phys. Lett. **265**, 115 (1997).
- ³² J. P. Perdew, M. Ernzerhof, and K. Burke, J. Chem. Phys. **105**, 9982 (1996).
- ³³ M. Ernzerhof, Chem. Phys. Lett. **263**, 499 (1996).
- ³⁴ P. J. Stephens, J. F. Devlin, and C. F. Chabalowski, J. Phys. Chem. **98**, 11623 (1994), (see also internet address <http://www.gaussian.com/q3.htm>).
- ³⁵ S. Raimes, *Many-Electron Theory* (North-Holland, Amsterdam, 1972).
- ³⁶ J. P. Finley, arXiv:physics/0506037 (2005).
- ³⁷ J. P. Finley, Phys. Rev. A (2005), (submitted).
- ³⁸ J. P. Finley, arXiv:physics/0506109 (2005).
- ³⁹ J. P. Finley, Phys. Rev. Lett. (2005), (submitted).
- ⁴⁰ J. P. Finley, Phys. Rev. A. **69**, 042514 (2004).
- ⁴¹ J. P. Finley, arXiv:physics/0308056 (2003).
- ⁴² J. P. Finley, Mol. Phys. **102**, 627 (2004).
- ⁴³ J. P. Finley (2005), (to be published).
- ⁴⁴ J.-P. Blaizot and G. Ripka, *Quantum Theory of Finite Systems* (MIT Press, Cambridge, Mass., 1986).
- ⁴⁵ J. P. Finley, Mol. Phys. **102**, 627 (2004), also arXiv:physics/0308084.
- ⁴⁶ R. M. Mattuck, *A Guide to Feynman Diagrams in the Many-Body Problem* (McGraw-Hill, New York, 1976), 2nd ed.